

PEEL FROM CORNER OF BOOKLET

15M-35-79H

Released For Shipment By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87 • BONHAM, TEXAS 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-Tx-001
Visit Us At: www.fertilome.com



STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

DANGER: Corrosive. Causes eye damage. Do not get in eyes or on skin or clothing. Wear goggles when applying this product. Harmful if swallowed or inhaled. Avoid exposure to spray mist. When using this product, wear long-sleeved shirt, long pants, socks, shoes, rubber gloves and eye protection. It is recommended that safety glasses include front, brow and temple protection. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.



ferti·lome



Weed-Out® Lawn Weed Killer
Contains **TRIMEC®** Herbicide

- **Kills troublesome broadleaf weeds: Dandelion, Chickweed, Clover and many others.**

ACTIVE INGREDIENTS:

Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid.....	5.45%
Dimethylamine salt of 2,4-dichlorophenoxyacetic acid.....	5.88%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid).....	1.21%
OTHER INGREDIENTS:	87.46%
TOTAL:	100.00%

THIS PRODUCT CONTAINS:
0.39 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 4.50%.
0.42 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 4.88%.
0.09 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 1.00%.
Isomer Specific by AOAC Methods.
Contains the single isomer form of Mecoprop-p.
TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**KEEP OUT OF REACH
OF CHILDREN
DANGER**

**SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS
AND FIRST AID
– KEEP FROM FREEZING-
NET CONTENTS 32 OZS.**

FIRST AID

If In Eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a Poison Control Center or doctor for treatment advice.
If On Skin Or Clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a Poison Control Center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none">• Call a Poison Control Center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a Poison Control Center or doctor.• Do not give anything by mouth to an unconscious person.
If Inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a Poison Control Center or doctor for further treatment advice.
Have the product container or label with you when calling a Poison Control Center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-800-858-7378 or 1-800-222-1222.	
Note To Physician: Probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. Store in original container and place in locked storage area when not using.
CONTAINER DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available. **If Partially Filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WHEN TO USE

Spray at any time when Weeds are growing. For best results, apply in Spring when Weeds are small and growing rapidly. Apply in Fall to remove late germinating Weeds.

HOW TO USE

The maximum application rate to turf is 0.6 pounds of 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Hose-End Sprayer: Use 1 gallon of water for every 250 square feet to be sprayed. For concentrate sprayers, add only the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart depending on type of Grass and number of square feet to be treated. Do not add water to concentrate sprayers. For dilution sprayers, add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** and fill sprayer with water to proper level. Spray uniformly.

Pressure-Sprayer: Use 1/2 gallon of water for every 250 square feet to be sprayed. Add the required amount of **ferti-lome® Weed Out Lawn Weed Killer** listed in the rate chart for the type of Grass to be treated and fill sprayer with water to the proper level. Adjust spray nozzle to deliver coarse spray. Spray uniformly.

Spot-Treatment: If hardy Weeds persist in Sensitive Grasses such as St. Augustine, Centipedegrass, Bermudagrass, Buffalograss or Bentgrass, spot treat individual Weeds using 1 teaspoon of **ferti-lome® Weed Out Lawn Weed Killer** in 1 quart of water (equivalent to 4 teaspoons per gallon of water.) Minimize contact of Grass.

RATES OF ferti-lome® Weed Out Lawn Weed Killer FOR VARIOUS LAWN GRASSES

Gallons Of Water Spray		Area to Treat (Sq. Ft.)	St. Augustinegrass, Centipedegrass (fl. oz.)	Bermudagrass, Buffalograss Bentgrass	Bluegrass Zoysiagrass, Fescue (fl. oz)
Hose End Sprayer	Pressure Sprayer				
1	1/2	250	1/2	2/3	2/3 to 1
4	2	1,000	2	2 2/3	2 2/3 to 4
8	4	2,000	4	5 1/3	5 1/3 to 8
20	10	5,000	10	13 1/3	13 1/3 to 20
32	16	8,000	16	21 1/3	21 1/3 to 32

EQUAL MEASURES

1 fluid ounce (fl. oz.) = 2 tablespoons (Tbs.) or 6 teaspoons (tsp.)
 2 fluid ounces = 4 tablespoons 4 fluid ounces = 1/2 cup 8 fluid ounces = 1 cup
 16 fluid ounces = 1 pint (pt.) 32 fluid ounces = 1 quart (qt.)

HOW TO USE IN THE 20 GALLON HOSE-END SPRAYER

1. Connect sprayer nozzle to garden hose; remove safety cap from container and insert sprayer unit, securing tightly.
2. Turn on water; place finger over hole on right side of nozzle and product will automatically be dispersed.
3. Spray evenly and uniformly over area requiring application. (Check application guide to determine the correct area to be covered per ounce.)

APPLICATION GUIDE

Fluid Ounces	Recommended Coverage Area Per Ounce		
	Sensitive Grasses Such As Centipedegrass And St. Augustinegrass (sq. ft.)	Bermudagrass, Bentgrass, And Buffalograss (sq. ft.)	Bluegrass, Fescue, And Zoysiagrass (sq. ft.)
1	500	375	250
2	1,000	750	500
3	1,500	1,125	750
4	2,000	1,500	1,000
5	2,500	1,875	1,250
6	3,000	2,250	1,500
7	3,500	2,525	1,750
8	4,000	3,000	2,000
9	4,500	3,375	2,250
10	5,000	3,750	2,500
11	5,500	4,125	2,750
12	6,000	4,500	3,000
13	6,500	4,875	3,250
14	7,000	5,250	3,500
15	7,500	5,625	3,750
16	8,000	6,000	4,000

Use Precautions: Do not apply this product through any type of irrigation system. Use only lawn sprayer that delivers coarse spray to reduce drift. Avoid spray drift to Vegetables, Flowers, Ornamental Plants. Do not spray roots of Trees and Ornamentals. Do not exceed specified dosages for any area; be particularly careful within dripline of Trees and

other Ornamental species. Do not spray Carpetgrass, Dichondra, or Lawn where desirable Clovers are present. Do not apply this product to "Floritam" St. Augustine Grass in Florida. Do not apply to newly seeded Lawn Grasses until well established. Do not spray when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to follow above precautions may result in injury.

GENERAL INFORMATION

Ferti-lome® Weed Out Lawn Weed Killer controls troublesome common Lawn Weeds, including Poison Ivy, in both Northern and Southern Grasses.

- BROADLEAF WEEDS -

Annual Yellow Sweetclover	Broadleaf Plantain	Clover, Crimson	Florida Pusley
Aster	Broomweed	Clover, Hop	Frenchweed
Austrian Fieldcress	Buckhorn	Clover, Red	Galinsoga
Bedstraw	Buckhorn Plantain	Clover, Strawberry	Goldenrod
Beggarticks	Bulbous Buttercup	Clover, Sweet	Ground Ivy
Betony, Florida	Bull Thistle	Clover, White	Gumweed
Bindweed, Field	Bullnettle	Cockle	Hairy Fleabane
Bird Vetch	Burclover	Cocklebur	Hawkweed
Bitter Wintercress	Burdock	Common Mullein	Healall
Bittercress, Hairy	Burning Nettle	Creeping Jenny	Heartleaf Drymary
Bitterweed	Burweed	Cudweed	Heathaster
Black-Eyed Susan	Buttercup	Curly Dock	Hedge Bindweed
Black Medic	Buttonweed	Daisy, English	Hedge Mustard
Black Mustard	Canada Thistle	Daisy Fleabane	Henbit
Blackseed Plantain	Carolina Geranium	Daisy, Oxeye	Hoary Cress
Blessed Thistle	Carpetweed	Dandelion	Hoary Plantain
Bloodflower Milkweed	Catchweed Bedstraw	Dichondra	Hoary Vervain
Blue Lettuce	Catnip	Dogfennel	Horsenettle
Blue Vervain	Catsear	Dollarweed	Jimsonweed
Bracted Plantain	Chickweed, Common	False Dandelion	Knawel
Bristly Oxtongue	Chickweed, Mouseear	Falseflax	Knotweed
Broadleaf Dock	Chicory	False Sunflower	Kochia
	Cinquefoil	Fiddleneck	Lambsquarters

- BROADLEAF WEEDS - (Continued)

Lespedeza	Poorjoe	Speedwell	Wild Carrot
Mallow	Prairie Sunflower	Spiny Amaranth	Wild Four-O'Clock
Matchweed	Prickly Lettuce	Spiny Cocklebur	Wild Garlic
Mexicanweed	Prickly Sida	Spiny Sowthistle	Wild Geranium
Milk Vetch	Prostrate Knotweed	Spotted Catsear	Wild Lettuce
Morningglory	Prostrate Pigweed	Spotted Spurge	Wild Marigold
Mouseear Hawkweed	Prostrate Spurge	Spurweed	Wild Mustard
Mugwort	Prostrate Vervain	Stinging Nettle	Wild Onion
Musk Thistle	Puncturevine	Strawberry, India Mock	Wild Parsnip
Mustard	Purslane, Common	Tall Nettle	Wild Radish
Narrowleaf Plantain	Ragweed	Tall Vervain	Wild Rape
Narrowleaf Vetch	Red Sorrel	Tansy Ragwort	Wild Strawberry
Nettle	Redroot Pigweed	Tansy Mustard	Wild Sweet Potato
Orange Hawkweed	Redstem Filaree	Tanweed	Wild Vetch
Oriental Cocklebur	Rough Cinquefoil	Thistle	Woodsorrel
Oxalis	Rough Fleabane	Trailing Crownvetch	Woolly Croton
Parsley-Piert	Russian Pigweed	Tumble Mustard	Woolly Morningglory
Parsnip	Russian Thistle	Tumble Pigweed	Woolly Plantain
Pearlwort	Scarlet Pimpernel	Velvetleaf	Wormseed
Pennycress	Scotch Thistle	Venice Mallow	Yarrow
Pennywort	Sheep Sorrel	Virginia Buttonweed	Yellow Rocket
Pepperweed	Sherherdspurse	Virginia Creeper	Yellowflower
Pigweed	Slender Plantain	Virginia Pepperweed	Pepperweed
Pineywoods Bedstraw	Smallflower Galinsoga	Wavyleaf Bullthistle	and other Broadleaf
Plains Coreopsis	Smooth Dock	Western Clematis	Weeds
Plantain	Smooth Pigweed	Western Salsify	
Poison Ivy	Sorrel	White Mustard	
Poison Oak	Sowthistle	Wild Aster	
Pokeweed	Spanishneedles	Wild Buckwheat	

LIMITED WARRANTY AND DISCLAIMER

The Manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This Limited Warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the Manufacturer or the Seller. Any damages arising from a breach of the Manufacturer's Warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss or profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the Seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.



Weed Out Lawn Weed Killer

Released For Shipment By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87 • BONHAM, TX 75418
EPA Reg. No. 2217-597-7401
EPA Est. No. 7401-TX-001
Visit Us At: www.fertilome.com

